

<p>(51) International Patent Classification ⁶ : A61K 39/00, 39/39, 48/00, C07K 14/47</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/20302</p> <p>(43) International Publication Date: 29 April 1999 (29.04.99)</p>
<p>(21) International Application Number: PCT/US98/22145</p> <p>(22) International Filing Date: 20 October 1998 (20.10.98)</p> <p>(30) Priority Data: 08/954,643 20 October 1997 (20.10.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/954,643 (CIP) Filed on 20 October 1997 (20.10.97)</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NI, SN, TD, TG).</p>
<p>(71) Applicant (for all designated States except US): AVANT IMMUNOTHERAPEUTICS, INC. [US/US]; 119 Fourth Avenue, Needham, MA 02194 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): RITTERSHAUS, Charles, W. [US/US]; 65 Garden Street, Malden, MA 02148 (US). THOMAS, Lawrence, J. [US/US]; 1 Fox Ridge Road, Easton, MA 02375 (US).</p>		<p>Published</p> <p>With international search report.</p> <p>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
<p>(74) Agent: YANKWICH, Leon, R.; Yankwich & Associates, 130 Bishop Allen Drive, Cambridge, MA 02139 (US).</p>		

Substance	% Inhibition
POSITIVE CONTROL MAB	100
HCG	~28
PEPTIDE	~72
rhUCETP	100

Methods for modulating cholesteryl ester transfer protein (CETP) activity and the plasma levels of lipoproteins involved in heart disease involve administration of a non-endogenous CETP or a plasmid-based vaccine for expression of such non-endogenous CETP to elicit production in a mammal of antibodies that recognize (bind to) the mammal's native (endogenous) CETP.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**XENOGENEIC CHOLESTERYL ESTER
TRANSFER PROTEIN (CETP) FOR MODULATION
OF CETP ACTIVITY**

5 **Cross-Reference to Related Application**

 This application is a continuation-in-part application of United States Patent Application Serial No. 08/954,643, filed 20 October 1997.

10 **Field of the Invention**

 This invention relates to the field of cardiovascular disease, particularly atherosclerosis. More particularly, the present invention provides compositions and methods for the control, treatment, or reduction of the risk of atherogenic activity in the circulatory system of mammals, particularly humans.

15 **Background of the Invention**

 Cholesterol circulates through the body predominantly as a component of lipoprotein particles (lipoproteins), which are composed of a protein portion, called apolipoproteins (Apo) and various lipids, including phospholipids, triglycerides, cholesterol and cholesteryl esters. There are ten major classes of apolipoproteins: Apo A-I, Apo A-II, Apo-IV, Apo B-20 48, Apo B-100, Apo C-I, Apo C-II, Apo C-III, Apo D, and Apo E. Lipoproteins are classified by density and composition. For example, high density lipoproteins (HDL), one function of which is to mediate transport of cholesterol from peripheral tissues to the liver, have a density usually in the range of approximately 1.063-1.21 g/ml. HDL contain various amounts of Apo A-I, Apo A-II, Apo C-I, Apo C-II, Apo C-III, Apo D, Apo E, as well as 25 various amounts of lipids, such as cholesterol, cholesteryl esters, phospholipids, and triglycerides (TG).

 In contrast to HDL, low density lipoproteins (LDL), which generally have a density of approximately 1.019-1.063 g/ml, contain Apo B-100 in association with various lipids. In particular, the amounts of the lipids, cholesterol, and cholesteryl esters are considerably 30 higher in LDL than in HDL when measured as a percentage of dry mass. LDL are particularly important in delivering cholesterol to peripheral tissues.

 Very low density lipoproteins (VLDL) have a density of approximately 0.95-1.006 g/ml and also differ in composition from other classes of lipoproteins both in their protein and lipid content. VLDL generally have a much higher amount of triglycerides than do HDL or

LDL and are particularly important in delivering endogenously synthesized triglycerides from liver to adipose and other tissues. The features and functions of various lipoproteins have been reviewed (see, for example, Mathews and van Holde, Biochemistry, pp. 574-576 and 626-630 (The Benjamin/Cummings Publishing Co., Redwood City, California, 1990); Havel et al., "Introduction: Structure and metabolism of plasma lipoproteins", in The Metabolic Basis of Inherited Disease, 6th ed., pp. 1129-1138 (Scriver et al., eds.) (McGraw-Hill, Inc., New York, 1989); Zannis et al., "Genetic mutations affecting human lipoproteins, their receptors, and their enzymes", in Advances in Human Genetics, Vol. 21, pp. 145-319 (Plenum Press, New York, 1993)).

Decreased susceptibility to cardiovascular disease, such as atherosclerosis, is generally correlated with increased absolute levels of circulating HDL, with lowered levels of LDL or VLDL, and also with increased levels of HDL relative to circulating levels of VLDL and LDL (see, e.g., Gordon et al., *N. Engl. J. Med.*, 321: 1311-1316 (1989); Castelli et al., *J. Am. Med. Assoc.*, 256: 2835-2838 (1986); Miller, et al., *Am. Heart J.*, 113: 589-597 (1987); Tall, *J. Clin. Invest.*, 89: 379-384 (1990); Tall, *J. Internal Med.*, 237: 5-12 (1995)).

Cholesteryl ester transport protein (CETP) mediates the transfer of cholesteryl esters from HDL to TG-rich lipoproteins such as VLDL and LDL, and also the reciprocal exchange of TG from VLDL to HDL (Tall, *ibid.*; Tall, *J. Lipid Res.*, 34: 1255-1274 (1993); Hesler et al., *J. Biol. Chem.*, 262: 2275-2282 (1987); Quig et al., *Ann. Rev. Nutr.*, 10: 169-193 (1990)). CETP may play a role in modulating the levels of cholesteryl esters and triglyceride associated with various classes of lipoproteins. A high CETP cholesteryl ester transfer activity has been correlated with increased levels of LDL-associated cholesterol and VLDL-associated cholesterol, which in turn are correlated with increased risk of cardiovascular disease (see, e.g., Tato et al., *Arterioscler. Thromb. Vascular Biol.*, 15: 112-120 (1995)).

CETP isolated from human plasma is a hydrophobic glycoprotein having 476 amino acids and a molecular weight of approximately 66,000 to 74,000 daltons on sodium dodecyl sulfate (SDS)-polyacrylamide gels (Albers et al., *Arteriosclerosis*, 4: 49-58 (1984); Hesler et al., *J. Biol. Chem.*, 262: 2275-2282 (1987); Jarnagin et al., *Proc. Natl. Acad. Sci. USA*, 84: 1854-1857 (1987)). A cDNA encoding human CETP has been cloned and sequenced (see, Drayna et al., *Nature*, 327: 632-634 (1987)). CETP has been shown to bind cholesteryl esters, triglycerides, phospholipids (Barter et al., *J. Lipid Res.*, 21:238-249 (1980)), and lipoproteins (see, e.g., Swenson et al., *J. Biol. Chem.*, 264: 14318-14326 (1989)). More recently, the

region of CETP defined by the carboxyl terminal 26 amino acids, and in particular amino acids 470 to 475, has been shown to be especially important for neutral lipid binding involved in neutral lipid transfer (Hesler et al., *J. Biol. Chem.*, 263: 5020-5023 (1988)).

Hereinafter, LDL-C will be used to refer to total cholesterol, including cholesteryl esters and/or unesterified cholesterol, associated with low density lipoprotein. VLDL-C will be used to refer to total cholesterol, including cholesteryl esters and/or unesterified cholesterol, associated with very low density lipoprotein. HDL-C will be used to refer to total cholesterol, including cholesteryl esters and/or unesterified cholesterol, associated with high density lipoprotein.

A number of *in vivo* studies utilizing animal models or humans have indicated that CETP activity can affect the level of circulating cholesterol-containing HDL. Increased CETP cholesteryl ester transfer activity can produce a decrease in HDL-C levels relative to LDL-C and/or VLDL-C levels which in turn is correlated with an increased susceptibility to atherosclerosis. Injection of partially purified human CETP into rats (which normally lack CETP activity), resulted in a shift of cholesteryl ester from HDL to VLDL, consistent with CETP-promoted transfer of cholesteryl ester from HDL to VLDL (Ha et al., *Biochim. Biophys. Acta*, 833: 203-211 (1985); Ha et al., *Comp. Biochem. Physiol.*, 83B: 463-466 (1986); Gavish et al., *J. Lipid Res.*, 28: 257-267 (1987)). Transgenic mice expressing human CETP were reported to exhibit a significant decrease in the level of cholesterol associated with HDL (see, e.g., Hayek et al., *J. Clin. Invest.*, 90: 505-510 (1992); Breslow et al., *Proc. Natl. Acad. Sci. USA*, 90: 8314-8318 (1993)). Furthermore, whereas wild-type mice are normally highly resistant to atherosclerosis (Breslow et al., *ibid.*), transgenic mice expressing a simian CETP were reported to have an altered distribution of cholesterol associated with lipoproteins, namely, elevated levels of LDL-C and VLDL-C and decreased levels of HDL-C (Marotti et al., *Nature*, 364: 73-75 (1993)). Transgenic mice expressing simian CETP also were more susceptible to dietary-induced severe atherosclerosis compared to non-expressing control mice and developed atherosclerotic lesions in their aortas significantly larger in area than those found in the control animals and having a large, focal appearance more typical of those found in atherosclerotic lesions in humans (Marotti et al., *ibid.*). Intravenous infusion of anti-human CETP monoclonal antibodies (Mab) into hamsters and rabbits inhibited CETP activity *in vivo* and resulted in significantly increased levels of HDL-C, decreased levels of HDL-triglyceride, and increased HDL size; again implicating a critical role for CETP in the

distribution of cholesterol in circulating lipoproteins (Gaynor et al., *Atherosclerosis*, 110: 101-109 (1994) (hamsters); Whitlock et al., *J. Clin. Invest.*, 84: 129-137 (1989) (rabbits)).

CETP deficiency has also been studied in humans. For example, in certain familial studies in Japan, siblings that were homozygous for non-functional alleles of the CETP gene had no detectable CETP activity. Virtually no atherosclerotic plaques were exhibited by these individuals, who also showed a trend toward longevity in their families (see, e.g., Brown et al., *Nature*, 342: 448-451 (1989); Inazu et al., *N. Engl. J. Med.*, 323: 1234-1238 (1990); Bisgaier et al., *J. Lipid Res.*, 32: 21-23 (1991)). Such homozygous CETP-deficient individuals also were shown to have an anti-atherogenic lipoprotein profile as evidenced by elevated levels of circulating HDL rich in cholesteryl ester, as well as overall elevated levels of HDL, and exceptionally large HDL, i.e., up to four to six times the size of normal HDL (Brown et al., *supra*, p. 451). The frequency of this mutation in Japan is relatively high, and may account for an elevated level of HDL in a significant fraction of the Japanese population.

The above studies indicate that CETP plays a major role in transferring cholesteryl ester from HDL to VLDL and LDL, and thereby in altering the relative profile of circulating lipoproteins to one which is associated with an increased risk of cardiovascular disease (e.g., decreased levels of HDL-C and increased levels of VLDL-C and LDL-C). Taken together, the current evidence suggests that increased levels of CETP activity may be predictive of increased risk of cardiovascular disease. Modulation or inhibition of endogenous CETP activity is thus an attractive therapeutic method for modulating the relative levels of lipoproteins to reduce or prevent the progression of, or to induce regression of, cardiovascular diseases, such as atherosclerosis.

In our previous work, detailed, e.g., in commonly assigned, copending patent application PCT/US96/06147 (WO 96/34888) and commonly assigned copending patent application PCT/US97/07294 (WO 97/41227), both incorporated herein by reference, we detailed an approach for modulating the CETP activity in an individual via vaccination with a peptide composition or with a plasmid-based vaccine that would lead to the production of antibodies recognizing and neutralizing endogenous CETP. We demonstrated that administration of immunogenic peptides either by direct inoculation or by *in situ* production following injection of a functional plasmid-based vaccine resulted in production of antibodies reactive with the inoculated individual's own (endogenous) CETP. Thus the vaccine peptides and the plasmid-based vaccines break tolerance in the vaccinated individuals and to promote

production of antibodies recognizing a self protein. Furthermore, administration of these vaccines to test animals resulted in a decline in the relative levels of total cholesterol and HDL-C and resulted in a decrease in the development of atherosclerotic lesions. The elicited endogenous anti-CETP antibodies thus promote a physiological condition correlated with decreased risk of cardiovascular disease, and they appear to have a direct effect on preventing or decreasing the formation of atherosclerotic plaques.

We have now discovered another approach to eliciting the production of anti-CETP antibodies in a mammal. We have now determined that whole CETP molecules of another mammal, that is, non-endogenous CETP, can be used to elicit antibodies in a mammal that will be reactive with its own, endogenous CETP and serve to modulate the activity of CETP and to provide lowered circulating CETP activity, lowered total cholesterol, lowered circulating LDL levels, elevated ratios of HDL-C to LDL-C. The use of non-endogenous CETP to promote production of anti-endogenous CETP antibodies also leads to a reduction in development of atherosclerotic lesions in comparison to unvaccinated controls.

Summary of the Invention

Accordingly, the present invention provides compositions and methods useful for the modulation or inhibition of cholesteryl ester transfer protein (CETP) activity. In particular, the use of non-endogenous CETPs, including xenogeneic CETPs, is described as a means, when administered to a mammal, to raise an antibody response against the mammal's own endogenous CETP and thereby to promote a prophylactic or therapeutic effect against cardiovascular disease, such as atherosclerosis. Such non-endogenous CETP can be CETP of another mammalian species (xenogeneic CETP), such as rabbit CETP, mouse CETP or simian CETP for administration to a human; the non-endogenous CETP can be a non-endogenous allelic variation or polymorph of a mammalian CETP administered to the same species of mammal (e.g., a human CETP polymorph administered to another human); or the non-endogenous CETP can be a CETP from one species modified to have an amino acid sequence more similar to the native CETP of another species (e.g., a "humanized" rabbit CETP for administration to a human).

Vaccine compositions and plasmid-based vaccines are described which, when suitably administered to a mammal result in the production of antibodies reactive with the mammal's endogenous CETP and the other benefits described herein.

The invention provides methods for eliciting antibodies in a mammal that will be reactive with its own, endogenous CETP, for modulating the activity of CETP in a mammal, and for providing in a mammal lowered circulating CETP activity, lowered total cholesterol, lowered circulating LDL levels, and/or elevated ratios of HDL-C to LDL-C. The invention provides a method for reducing or preventing in a mammal the development of atherosclerotic lesions.

Brief Description of the Drawings

Figure 1A-C. An alignment of the amino acid sequences of mature rabbit CETP (SEQ ID NO: 3) with mature human CETP (SEQ ID NO: 1). The rabbit CETP is shown over the aligned human CETP sequence. The rabbit sequence includes 20 more amino acid residues than the human sequence, and the human sequence shows a 1-amino acid and a 19-amino acid gap (indicated with dashes, --, in the human sequence) in order to show the residue matches (indicated with a vertical line, |) most clearly.

Figure 2. Shows the end point titers of antibodies from rabbit plasma recognizing the C-terminal peptide of rabbit CETP from rabbits vaccinated with human chorionic gonadotropin ("HCG Vaccine"), a synthetic vaccine peptide having segments of tetanus toxoid and the C-terminal sequence of human CETP ("Peptide Vaccine", see SEQ ID NO: 7), and full-length recombinant human CETP ("rhuCETP"). The figure shows maximum anti-CETP antibody titers achieved for each rabbit in an ELISA detecting plasma antibodies specific for rabbit CETP C-terminal peptide (amino acids 477-496).

Figure 3. Shows inhibition of CETP activity in a commercial fluorescence-based assay (Roar Biomedical, Yonkers, New York) by protein A-isolated antibodies from the plasma of vaccinated rabbits from Groups I-III (see Examples, *infra*). The graph shows percent inhibition achieved in each of the vaccinated rabbits.

Figure 4. Shows the CETP activity in the vaccinated rabbits from week 1 to week 32.

Figure 5. Shows the percentage change in total cholesterol levels in the vaccinated rabbits from week 1 to week 12.

Figure 6. Shows the HDL-associated cholesterol levels in the vaccinated rabbits from week 1 to week 32.

Figure 7. Shows the percentage change in LDL-associated cholesterol levels in the vaccinated rabbits from week 1 to week 12.

Figure 8. Shows the plasma lipoprotein levels for a rabbit vaccinated with non-endogenous CETP(recombinant human CETP, or rhuCETP). "V" indicates the periodic vaccination boosts. HDL-associated cholesterol, total cholesterol level, and triglyceride level were assayed.

5 Figure 9. Shows a correlation between CETP activity and HDL as a percent of total lipoproteins and total CETP mass, in a rabbit vaccinated with non-endogenous CETP (rhuCETP)

10 Figure 10. Shows the levels of cholesterol deposits in the irises of 48 rabbits vaccinated with human chorionic gonadotropin ("HCG", rabbits #1-#12), a synthetic vaccine peptide having segments of tetanus toxoid and the C-terminal sequence of human CETP ("Peptide", see SEQ ID NO: 7, rabbits #13-#24), full-length recombinant human CETP ("rhuCETP", rabbits #25-#36), and a CETP-tetanus toxoid conjugate composition. ("Conjugate", rabbits #37-#48). The irises of each rabbit have been scored based on the percentage of deposits observed from 0-5, 0 representing no deposits and 5
15 representing 100% deposits observed in the iris of the animal.

Figure 11. Shows percentage of the aorta covered by lesions observed in vaccinated rabbits fed an atherogenic diet. Values of individual animals are represented by open symbols.

20 Figure 12. Shows anti-CETP antibody titers for mice vaccinated with various plasmid-based vaccines.

Figure 13A-K. Shows antibody titers of 11 rabbits vaccinated with pSV40-HuCETP between weeks 1-32. The open diamond and open square symbols refer to the antibody titers of the rabbits on weeks 30 and 34 respectively.

25 Figure 14. Shows the effect of plasmid vaccination on the development of aortic lesions. The figure compares the mean percentages collected for control rabbits (pSV40 plasmid without the CETP coding sequence) and rabbits vaccinated with pSV40-huCETP.

Detailed Description of the Invention

30 As noted above, a decreased risk of atherosclerosis has been correlated with relatively low circulating levels of LDL and VLDL and relatively high levels of HDL. The levels of such circulating lipoproteins are directly influenced, at least in part, by the endogenous levels of CETP activity. In particular, high CETP activity promotes transfer of neutral lipids, such

as cholesteryl esters from HDL to VLDL and LDL. Accordingly, CETP is a relatively precise target in humans and other animals for altering the relative levels of LDL, VLDL and HDL in the circulatory system (see, e.g., Tato et al., *Arteriosclero. Thromb. Vascular Biol.*, 15: 112-120 (1995); Tall, *J. Internal Med.*, 237: 5-12 (1995)). This invention is directed to the control of endogenous CETP activity by providing non-endogenous CETP molecules to an individual, for promoting an immune response in such individuals against their endogenous CETP, thereby promoting a physiological condition (e.g., increased level of HDL or decreased level of LDL) correlated with a decreased risk of atherosclerosis. In addition, promoting an immune response against endogenous CETP using the vaccine compositions of this invention can provide, prevent, or inhibit the progression of lesions in tissue susceptible to atherosclerosis.

Compositions for Modulation of CETP Activity

As used herein, a CETP vaccine composition for use according to the invention contains as an essential ingredient a CETP or a portion thereof, that is non-endogenous with respect to the mammal to be vaccinated. For the purposes of this invention, "non-endogenous CETP" means a cholesteryl ester transfer protein that is not the native CETP produced by the mammal to be vaccinated. For example, with respect to a human subject, non-endogenous CETP will include CETP produced by another mammalian species, i.e., xenogeneic CETP, such as rabbit, mouse or simian CETP; or non-endogenous CETP with respect to a particular human subject can be an allelic variant or polymorphism of human CETP, such as CETP produced by another human individual.

The non-endogenous CETP can also be a xenogeneic CETP that has been modified in order to make the amino acid sequence of the modified CETP more similar to that of the endogenous CETP of the mammal to be vaccinated. The term used herein to describe such modified non-endogenous CETP is "mammalianized CETP". This term is used with reference to the mammal to be vaccinated, and it means a non-endogenous CETP that has been modified to have an amino acid content more similar to the native CETP of said mammal. An example of a mammalianized CETP, where the subject to be vaccinated is a human, would be a rabbit CETP modified (or "humanized") to have an amino acid sequence more similar to the native human sequence. As a further example, reference is made to Figures 1A, 1B and 1C, which show the respective amino acid sequences of rabbit CETP (SEQ ID NO: 3) and human CETP (SEQ ID NO: 1) in alignment.

Referring to Figures 1A-1C, it is seen that the structure of these two mammalian CETPs is similar, having the same amino acids at 80% of the positions of human CETP. Rabbit CETP (SEQ ID NO: 3) is 20 amino acids longer than human CETP (SEQ ID NO: 1), and the alignment of the two proteins in Figures 1A-1C shows two segments, denoted with dashes (---), where the proteins do not correspond structurally. With respect to a human subject to be administered a non-endogenous CETP in accordance with this invention, an example of a "mammalianized" non-endogenous CETP would be a rabbit CETP in which the 19-amino acid segment from amino acid Ala₃₉₃ through Ala₄₁₁ of rabbit CETP has been deleted, making the modified CETP (see SEQ ID NO: 5) 477 amino acids in length and thus more similar to the human CETP (SEQ ID NO: 1). Since the mammal of this example is a human, another term for such a modified CETP or mammalianized non-endogenous CETP would be a "humanized rabbit CETP".

Again referring to Figures 1A-1C, a further example of a humanized rabbit CETP would be a CETP as set forth in SEQ ID NO: 6. In Figure 1C it is noted that in the C-terminal portion of the human and rabbit CETPs there is only one difference in the respective amino acid sequences, i.e., Lys₄₈₅ of the rabbit CETP corresponds to Glu₄₆₅ in the human CETP.

In practicing the methods of the present invention, non-endogenous CETP is administered to a mammal in an amount effective to elicit an immune response. As is common in the field, more than one administration may be necessary or desirable to obtain a high enough concentration of anti-endogenous CETP antibodies in the mammal to affect endogenous CETP activity.

Immunogenicity of a vaccine peptide of this invention may be further enhanced by linking the immunogenic non-endogenous CETP to itself or to a related protein homologous to CETP. In this approach, a dimer could be formed, with the dimer providing a multi-chain protein that is even more immunogenic than the non-endogenous CETP alone. Examples of proteins related to CETP that might be used in this approach include, for example, phospholipid transfer protein and neutrophil bactericidal protein (see, Day et al., *J. Biol. Chem.*, 269: 9388-91 (1994); Gray et al., *J. Biol. Chem.*, 264: 9505-9509 (1989)).

Other immunogenic carrier molecules such as keyhole limpet hemocyanin (KLH) may also be used in combination with the non-endogenous CETP. For example, KLH contains multiple lysine residues in its amino acid sequence. Each of these lysines is a potential site at

which a CETP molecule as described herein could be linked (for example, using maleimide-activated KLH, Catalog No. 77106, Pierce, Rockford, IL), to provide a multivalent non-endogenous CETP immunogen. Another example of an immunogenic carrier molecule is hsp70 from *Mycobacterium tuberculosis*, which has been shown to be an especially potent antigen containing multiple B and T cell epitopes (see, e.g., Suzue and Young, *J. Immunol.*, 156: 873 - 879 (1996)). The hsp70 protein can be linked by standard cross-linking agents to non-endogenous CETPs to enhance immunogenicity of the vaccine compositions.

Other peptides can be conjugated with the non-endogenous CETP molecules to provide a source of helper T cell epitopes and boost the immunogenicity of the vaccine compositions according to the invention. Such peptides include, for example, "universal" antigenic peptides, e.g., tetanus toxoid or diphtheria toxoid, especially the tetanus toxoid fragment: Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu (amino acids 2 to 15 of SEQ ID NO: 7).

Pharmaceutically acceptable adjuvants, such as alum, may be mixed with the non-endogenous CETPs described herein to produce vaccine compositions of this invention. Alum is the single adjuvant currently approved for use in administering vaccines to humans (see, Eldridge et al., in Immunobiology of Proteins and Peptides V: Vaccines: Mechanisms, Design, and Applications (Atassi, M.Z., ed.) (Plenum Press, New York, 1989), page 192). Alum in combination with a sodium phthalyl derivative of lipopolysaccharide can also be used (see, Talwar et al., *Proc. Natl. Acad. Sci. USA*, 91: 8532-8536 (1994)). Other conventional adjuvants may be used as they are approved for a particular use. For example, biodegradable microspheres comprised of poly (DL-lactide-co-glycolide) (Eldridge et al., *supra*, pp. 191-202); Freund's Complete Adjuvant (Sigma Chemical Co., St. Louis, Missouri), Freund's Incomplete Adjuvant (Sigma Chemical Co., St. Louis, Missouri), and the RIBI™ Adjuvant System (RAS; RIBI ImmunoChem Research, Inc., Hamilton, Montana); lipophilic N-palmitoyl-S-[2,3-bis(palmitoyloxy)-propyl]-cysteine ("Pam₃-Cys-OH"); amphiphilic, water-soluble lipopeptides such as Pam₃-Cys-Ser-Lys, and Pam₃-Cys-Ser-Glu₄; glycopeptides such as N-acetyl-glucosaminyl-N-acetylmuramyl-alanyl-D-isoglutamine ("GMDP"), muramyl dipeptides, and alanyl-N-adamantyl-D-glutamine; and polyamide gel-based adjuvants which can easily be attached to CETP peptides during their *in vitro* chemical synthesis (see, Synthetic Vaccines (Nicholson, ed.) (Blackwell Scientific Publications, Cambridge, Massachusetts, 1994), pp. 236-238).

Where helper T cell epitope molecules or adjuvant species are to be physically linked or conjugated with the non-endogenous CETP, the CETP can be covalently linked directly or via a cross-linker molecule.

Suitable cross-linking molecules include amino acids, for example, using one or more glycine residues to form a "glycine bridge" between the CETP and the carrier or adjuvant molecule. Also contemplated are the formation of disulfide bonds between cysteine residues, or the use of cross-linking molecules such as glutaraldehyde (see, Korn et al., *J. Mol. Biol.*, 65: 525-529 (1972)) and EDC (Pierce, Rockford, IL) or other bifunctional cross-linker molecules. Bifunctional cross-linker molecules possess two distinct reactive sites; one of the reactive sites can be reacted with a functional group on the CETP and the other reactive site can be reacted with a functional group on the carrier or adjuvant molecule, uniting the two. General methods for cross-linking molecules are reviewed by Means and Feeney (*Bioconjugate Chem.*, 1: 2-12 (1990)).

Homobifunctional cross-linker molecules have two reactive sites which are chemically the same. Examples of homobifunctional cross-linker molecules include glutaraldehyde; N,N'-bis(3-maleimido-propionyl)-2-hydroxy-1,3-propanediol (a sulfhydryl-specific homobifunctional cross-linker); certain N-succinimide esters, such as disuccinimidyl suberate and dithio-bis-(succinimidyl propionate) and their soluble bis-sulfonic acids and salts (e.g., as available from Pierce Chemicals, Rockford, Illinois; or Sigma Chemical Co., St. Louis, Missouri).

Preferably, the bifunctional cross-linker molecule is a heterobifunctional linker molecule, meaning that the linker molecule has at least two reactive sites that can be separately covalently attached to the non-endogenous CETP and the carrier or adjuvant molecule. Heterobifunctional cross-linker molecules that may be used include m-maleimidobenzoyl-N-hydroxysuccinimide ester; m-maleimido-benzoylsulfosuccinimide ester; γ -maleimidobutyric acid N-hydroxysuccinimide ester; and N-succinimidyl 3-(2-pyridyl-dithio)propionate.

The non-endogenous CETP for use according to this invention can be produced by any of the available methods known in the art to produce proteins of defined amino acid sequence. For example, automated peptide synthesis is available to those skilled in the art by using automated peptide synthesizers (e.g., Synergy™ Peptide Synthesizer by Applied Biosystems; AMS 422 by Abimed, Langenfeld, Germany). Synthesis of such proteins to

order is performed as a commercial service by many commercial peptide synthesizing service companies, e.g., Quality Controlled Biochemicals, Inc., (Hopkinton, Massachusetts); Chiron Mimotopes Peptide Systems (San Diego, California); Bachem Bioscience, Inc. (Philadelphia, Pennsylvania); Severn Biotech Ltd. (Kiddeminster, England).

5 Alternatively, the proteins of this invention may be produced using synthetic and recombinant nucleic acid technology. For example, one of ordinary skill in the art can design from the known genetic code a 5' to 3' nucleic acid sequence encoding a proteins of this invention. The amino acid sequence for a mature human CETP is known (SEQ ID NO: 1), as well as its corresponding DNA sequence (SEQ ID NO: 2) (see, Drayna et al., *Nature*, 327: 10 632 - 634 (1987)). Furthermore, the amino acid sequence for a mature rabbit CETP is known (SEQ ID NO: 3), as well as its corresponding DNA sequence (SEQ ID NO: 4) (see, Nagashima et al., *J. Lipid. Res.*, 29: 1643 - 1649 (1988)). A DNA molecule containing the coding sequences of desired CETP (or a modified "mammalianized" CETP as described above) can readily be synthesized either using an automated DNA synthesizer (e.g., Oligo 15 1000 DNA Synthesizer, Beckman Corp.) or by contracting with a commercial DNA synthesizing service (e.g., Genset Corp., La Jolla, California).

 The synthesized or cloned DNA molecule can then be inserted into any of a variety of available gene expression systems (e.g., bacterial plasmids; bacteriophage expression vectors, retroviral expression vectors, baculoviral expression vectors), using standard methods 20 available in the art (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, Vols. 1-3 (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989)) and/or as directed by the manufacturer of a particular commercially available gene expression system (e.g., pPROEX™-1 bacterial cell expression system; SFV eukaryotic cell expression system; CHO (Chinese Hamster Ovary cells) expression system; BAC-TO-BAC™ baculovirus expression 25 system; Life Technologies, Inc., Gaithersburg, Maryland). Especially preferred for the expression of CETP is the CHO expression system. The expressed CETP can then be isolated from the expression system using standard methods to purify proteins.

 Purification of the non-endogenous CETPs of this invention may be expedited by employing affinity chromatography or immunoprecipitation based on using antibodies to the 30 particular CETP domain. For example, the Mab TP2 (Hesler et al., *J. Biol. Chem.*, 263: 5020-5023 (1988)) binds to the carboxyl terminal 26 amino acids of human CETP, and could be useful in one or more immunoaffinity steps in a purification scheme. Another method that

could be used in the purification of the proteins is standard column chromatography (Weinberg et al., *J. Biol. Chem.*, 269: 29588-29591 (1994)).

The non-endogenous CETPs as described herein are used to make vaccines compositions that elicit, when administered to a mammal, production of endogenous antibodies which specifically bind to endogenous CETP of the mammal and/or modulate (i.e., decrease or inhibit) endogenous CETP activity in the mammal. The anti-CETP vaccine compositions of this invention may contain one or several different non-endogenous CETPs.

In addition, the non-endogenous CETP may be linked to other molecules that may enhance the immunogenicity of the peptides.

The vaccine compositions for administration of non-endogenous CETP according to this invention can also advantageously take the form of plasmid-based vaccines for producing non-endogenous CETP *in situ*, eliciting autoantibodies directed to endogenous CETP. Such plasmid-based vaccines are, specifically, DNA plasmids which are administered (for example, by intramuscular injection or intradermal ballistic administration) to an individual. The administered DNA plasmids encode and direct the production of immunogenic non-endogenous CETP. We have discovered that such immunogenic CETP elicits the production of autoantibodies that react specifically with (i.e., bind to) endogenous CETP in the individual.

An example of a recombinant plasmid that can be used to produce a non-endogenous CETP for use according to this invention is plasmid pCMV-CETP/TT in which the CMV promoter directs transcription of a sequence encoding a vaccine peptide described in the previously mentioned PCT/US96/06147. *E. coli* bearing plasmid pCMV-CETP/TT has been deposited with the American Type Culture Collection (ATCC, Rockville, MD) and assigned Accession No. 98038. DNA coding for a desired CETP can be inserted in place of the vaccine peptide coding sequence in pCMV-CETP/TT and used for the expression of full-length CETP molecules.

The production of anti-CETP antibodies promotes a physiological state associated with a decreased risk of cardiovascular disease. The beneficial modulation of CETP activity produced by the DNA vaccines is evidenced by a significantly decreased or eliminated CETP activity; by an anti-atherogenic lipoprotein profile (for example, an increase in the level of HDL or HDL-C compared to LDL, LDL-C, VLDL, or VLDL-C); or by an inhibition

(including prevention) or decrease in the development of atherosclerotic lesions in cardiovascular tissue, such as the aorta.

General methods of administering and testing vaccines are well known to those skilled in the art (see, e.g., Talwar et al., *Proc. Natl. Acad. Sci. USA*, 91: 8532-8536 (1994)). The immune response to endogenous CETP should significantly inhibit CETP activity, alter the serum half-life of CETP, cause clearance CETP through formation of immune complexes, alter the trafficking of HDL-cholesterol, shift the equilibrium of cholesterol content of lipoproteins, alter cholesterol catabolism, and/or reduce development of atherosclerotic lesions. Control of LDL, VLDL and/or HDL levels is an accepted indicator or endpoint in treatment of cardiovascular disease, as these levels are correlated with a decreased risk of cardiovascular disease or further progression of such disease (see, e.g., Mader, in Human Biology, 4th ed., pp. 83, 102 (Wm. C. Brown Publishers, Dubuque, Iowa, 1995)). Accordingly, the desired prophylactic or therapeutic effect according to this invention is evidenced by eliciting antibodies in an individual that bind to endogenous CETP and/or inhibit endogenous CETP activity, or by a relative decrease in LDL and/or VLDL levels compared to HDL levels as the titer of antibody directed against the endogenous CETP rises, or by a decrease of absolute levels of circulating LDL and/or VLDL with the production of anti-CETP antibodies, or by an inhibition or decrease in development of atherosclerotic lesions in cardiovascular tissue.

As demonstrated herein, administration of non-endogenous CETP in a rabbit model of atherosclerosis led to a significant decrease in the development of atherosclerotic plaques. This evidence indicates that vaccination to elicit antibodies to endogenous CETP may be a useful method of treating or preventing atherosclerosis.

The successful use of non-endogenous CETP to elicit anti-endogenous CETP antibodies and to modulate the activity of native CETP was surprising in a number of respects. Previously, the use of whole CETP molecules had been avoided, since it was not known whether introduction of a whole, non-endogenous CETP molecule would provide the desired immunogenic effects. For example, non-endogenous CETP might function perfectly well as a CETP and exacerbate already undesirable cholesterol levels and metabolism. In addition, it was contemplated that full-length CETP molecules might contain immunogenic segments that would elicit antibodies capable of reacting or interfering with proteins or receptors outside of the CETP metabolic pathway, resulting in dangerous cross-reactions or

side-effects. Finally, it was not known whether introduction of a non-endogenous CETP would be able to break tolerance in the subject vaccinated, leading to production of antibodies reactive not with (or not *only* with) the non-endogenous CETP but with the *native* CETP. These uncertainties have now been resolved.

5 The non-endogenous CETP vaccine compositions may be administered by any route used for vaccination, including: parenterally such as intraperitoneally, interperitoneally, intradermally, subcutaneously, intramuscularly, intravenously or orally. Preferably, the vaccines of this invention are administered parenterally, e.g., intraperitoneally, interperitoneally, subcutaneously, intradermally, intramuscularly, or intravenously. If oral
10 administration of a vaccine peptide is desired, any pharmaceutically acceptable oral excipient may be mixed with the vaccine peptides of this invention, for example, such as solutions approved for use in the oral polio vaccine. As with certain other vaccines, such as for tetanus, an occasional booster administration of the CETP vaccine compositions may be necessary to produce or maintain a desired level of modulation or inhibition of endogenous CETP.
15 Biodegradable microspheres, such as those comprised of poly (DL-lactide-co-glycolide), have been shown to be useful for effective vaccine delivery and immunization via oral or parenteral routes.

 Appropriate dosages of the non-endogenous CETP may be established by general vaccine methodologies used in the art based on measurable parameters for which the vaccine
20 is proposed to affect, including monitoring for potential contraindications, such as hypersensitivity reaction, erythema, induration, tenderness (see, e.g., Physician's Desk Reference, 49th ed., (Medical Economics Data Production Co., Mont Vale, New Jersey, 1995), pp. 1628, 2371 (referring to hepatitis B vaccine), pp. 1501, 1573, 1575 (referring to measles, mumps, and/or rubella vaccines), pp. 904, 919, 1247, 1257, 1289, 1293, 2363
25 (referring to diphtheria, tetanus and/or pertussis vaccines)) ; Talwar, G.P., et al., *Proc. Natl. Acad. Sci. USA*, 91: 8532-8536 (1994)). A common and traditional approach for vaccinating humans is to administer an initial dose of a particular vaccine to sensitize the immune system and then follow-up by one or more "booster" doses of the vaccine to elicit an anamnestic response by the immune system that was sensitized by the initial administration of the
30 vaccine (vaccination). Such a "primary and booster" administration procedure has been well known and commonly used in the art, as for example, in developing and using measles, polio, tetanus, diphtheria, and hepatitis B vaccines.

Initially, the amount of a vaccine composition administered to an individual may be that required to neutralize the approximate level of endogenous CETP activity present in the individual prior to vaccination, as can be determined by measuring CETP activity in serum or plasma samples from the individual, for example as determined using a commercially available CETP assay (e.g., Roar Biomedical., Yonkers, New York). Plasma or serum samples from a vaccinated individual can also be monitored to determine whether a measurable increase in the levels of total HDL or HDL-C is seen after administration of the non-endogenous CETP using commercially available assays (e.g., available from Sigma Diagnostics, Inc., Saint Louis, Missouri). A rise in the concentration (titer) of circulating anti-CETP antibodies can be measured in plasma or serum samples, for example using an ELISA assay. Thus, it is possible and recommended that initially it be established whether a rise in anti-CETP antibody can be correlated with an increase in the level of HDL or HDL-C, a decrease in LDL or VLDL, or with a decrease in CETP activity. Thereafter, one need only monitor a rise in titer of anti-CETP antibody to determine whether a sufficient dosage of vaccine peptide has been administered or whether a "booster" dose is indicated to elicit an elevated level of anti-CETP antibody. This is the common procedure with various established vaccinations, such as vaccination against hepatitis B virus.

Three-dimensional arterial imaging methods are currently available which can be used to identify arterial lesions and monitor their development or regression in an individual (see, for example, McPherson, *Scientific American Science & Medicine*, pages 22-31, (March/April, 1996)). Thus such imaging methods can be used to monitor the effectiveness of vaccination according to the methods of this invention.

A more complete appreciation of this invention and the advantages thereof can be obtained from the following non-limiting examples.

EXAMPLE 1

Immunization of Rabbits Against Endogenous CETP

Four vaccine preparations were made for injection into four groups of twelve New Zealand White Rabbits, to test the ability of the vaccine preparation to elicit an immune response against endogenous rabbit CETP. Group I (negative control) contained rabbits #1 - #12, each of which was injected with a vaccine composition containing an irrelevant antigen, human chorionic gonadotropin (hCG). Group II (comparative embodiment of

PCT/US96/06147) contained rabbits #13 - #24, each of which was injected with a vaccine peptide having a portion of the C-terminus of human CETP and a portion of tetanus toxoid ("Peptide"; see, SEQ ID NO: 7). Group III (this invention) contained rabbits #25 - #36, each of which received a vaccine composition containing whole recombinant human CETP ("rhuCETP"). Group IV (this invention) contained rabbits #37 - #48, each of which received a vaccine composition containing whole recombinant human CETP conjugated with whole tetanus toxoid using a chemical crosslinker ("Conjugate").

The general protocol for vaccinating and testing the rabbits was as follows: On Day 1, each rabbit received one subcutaneous injection of a composition containing 200 µg of immunogen in Complete Freund's Adjuvant (Sigma Chemical Co., St. Louis, Missouri). Each composition suspended the respective immunogen in phosphate buffered saline (PBS) and emulsified with complete Freund's adjuvant (1:1) to yield a final concentration of 100 µg/100 µl. Each rabbit was administered the vaccine mixture in a 200 µl dose (200 µg immunogen) at one subcutaneous site. A boost of 200 µg of immunogen in Incomplete Freund's Adjuvant (Sigma Chemical Co.) was administered as on Day 1 at Day 28 and Day 49. Blood samples (approximately 1-5 ml) were withdrawn from the ear of each rabbit (prior to injections) on Days 1 ("prebleed"), 28, 49, 105, 147 and 217. Blood plasma samples were prepared by standard centrifugation methods to separate cellular components from the plasma. Plasma samples were stored at -70° C. Plasma samples of both Groups I and II were analyzed for presence of and increase in titer of anti-CETP antibodies and for CETP activity, CETP mass, and plasma levels of various lipoprotein components (HDL, LDL, triglycerides).
Direct ELISA for Titering Anti-CETP Antibodies

A sandwich enzyme-linked immunosorbent assay (ELISA) was used to titer plasma samples containing anti-CETP antibody. A biotinylated C-terminal peptide (20 amino acids) of rabbit CETP was adsorbed to wells of a microtiter dish coated with streptavidin, and various dilutions of rabbit plasma from the rabbits of Groups I - III were added to each well. Non-specific binding can be blocked by adding a 1% solution of BSA in PBS and 0.05% Tween to each well and incubating for 2 hours at room temperature (20 -22° C) on a rotating shaker at 150 rpm. The wells were then washed four times with ELISA wash buffer (PBS + 0.05% Tween). Plasma samples were then diluted in dilution buffer (e.g. 1% BSA in PBS), followed by serial dilutions in the same buffer. Diluted samples (100 µl) were added to the wells, incubated for 1.5-2 hours at room temperature on a rotating shaker at 150 rpm, and

then washed 4 times with ELISA wash buffer (PBS + 0.05% Tween). To detect bound anti-CETP antibodies, 100 µl of an optimized dilution of horseradish peroxidase (HRP) labeled goat anti-rabbit immunoglobulin (Southern Biotechnology Associates, Inc.; Birmingham, Alabama; or Jackson ImmunoResearch, Inc.; West Grove, Pennsylvania) in dilution buffer was added, and the plates were incubated for 2 hours at room temperature on a rotating shaker at 150 rpm. The wells were then washed four times with ELISA wash buffer (see above), peroxidase substrate TMB (TMB peroxidase substrate, Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Maryland) was added, and the plates were incubated 30 minutes at room temperature. Change in optical density was monitored spectrophotometrically at 450 nm using an ELISA reader (e.g., E-max, Molecular Device Corp., Menlo Park, California). In this assay, the O.D. was directly proportional to the amount of anti-CETP antibodies present in the plasma samples.

The results of the assay for each of the rabbit groups is shown in Figure 2. Group I showed no plasma antibodies detecting the rabbit C-terminal CETP peptide, whereas the groups vaccinated with CETP immunogens (Groups II, III) showed significant titers of anti-CETP antibodies in almost all vaccinated rabbits. Analysis of the Group IV data is in progress.

CETP Activity and Neutralization Assays

In order to measure the activity of CETP in plasma, a commercial fluorescence-based assay (Roar Biomedical Inc.; Yonkers, New York) was used. Incubation of a CETP source (rhoCETP, huCETP or rabbit CETP) with the donor and acceptor particles, included in the kit, results in the CETP-mediated transfer of a fluorescent neutral lipid. This fluorescent neutral lipid is present in a self-quenched state when contained within the core of the donor. The CETP-mediated transfer is determined by the increase in fluorescence intensity as the fluorescent neutral lipid is removed from the self-quenched donor to the acceptor. To measure neutralization, anti-CETP antibodies are isolated from the plasma of vaccinated rabbits with protein A. Identical amounts (measured by A₂₈₀) of the antibodies from various samples are added to the above reaction.

Figure 3 shows inhibition of CETP activity by rabbit antibodies collected from the plasma of the vaccinated rabbits. The bar designated "Positive Control Mab" refers to anti-CETP monoclonal antibody TP2, which is known to inhibit CETP activity in vitro and is

included for comparison. Figure 4 shows the change in CETP activity in Groups I and III from week 1 to week 32. Analysis of the Group IV data is in progress.

Cholesterol and HDL Levels in Plasma Samples of Vaccinated Rabbits

The plasma samples taken from rabbits of Groups I - IV were also assayed for the concentration of total cholesterol (Figure 5), HDL-C (Figure 6), and triglycerides (see Figure 8) using standard commercial assays (Sigma Diagnostics, Inc., Saint Louis, Missouri). LDL-C (Figure 7) is calculated as total cholesterol minus HDL-C minus $0.2 \times$ tryglyceride level. CETP levels were determined by a slot blot analysis using anti-CETP monoclonal antibody TP2 and chemiluminescences for detection. The band intensities obtained with various amounts of plasma samples were quantified with the aid of a Kodak® DC40 camera and 1D Image Analysis software (version 1.6), then compared to that obtained with known amounts of purified human CETP loaded on the same nitrocellulose filter.

The plasma lipoprotein profile for the Group III rabbit #32, which showed the most pronounced reduction in LDL-C levels (Fig. 7) is shown in Figure 8. The profile shows a dramatic rise in HDL as a percent of the total lipoprotein profile. Figure 9 shows, for this rabbit, the correlation between decreasing CETP activity, decreasing cholesterol mass and increasing HDL as a percentage of total lipoprotein.

Measurement of cholesterol deposits in the irises of vaccinated rabbits

The rabbits from Groups I - IV were also assayed for the amount of cholesterol deposits detected in the irises. A scale of cholesterol deposition in the iris was established, with 0 = no deposit, 1 = 20%, 2 = 40%, 3 = 60%, 4 = 80%, and 5 = 100% deposits on this iris (i.e., iris completely covered with deposits). One iris per rabbit was evaluated and scored for degree of cholesterol deposition. The groups of animals were blinded to the scorer to avoid bias. Figure 10 shows the data collected from all 48 animals. The Xs indicate animals for which no data were obtained (rabbit #34, #39, and #42); these animals were euthanized due to unrelated complications such as furballs. The data indicate that all CETP vaccinated groups had statistically less cholesterol deposits than the control group.

Quantitation of lesions in aorta of vaccinated rabbits

The rabbits were switched from a diet of basic rabbit chow to diets supplemented with 0.25% cholesterol (w/w) known to produce atherosclerotic-like lesions in rabbits (Daley et al., *Arterioscler. Thromb.*, 14: 95 - 104 (1994)). To determine whether the vaccination may affect the development of atherosclerosis, the aortas of these rabbits were examined for the

development of atherosclerotic lesions. After blood samples were taken on the last day, rabbits were sacrificed. The entire aortas from each of Groups I-IV were removed and placed into fixative solution (3.7 % v/v formaldehyde). Loose tissue, adherent fat, and the adventitia were dissected free from the arteries. Each artery was then cut lengthwise, pinned flat to expose the intimal (luminal) surface, stained with Sudan IV, and then photographed. Sudan IV is a fat soluble red dye that stains atherosclerotic plaques on the intimal surface of arteries. Figure 11 summarizes the results of this experiment. The stained aortas of rabbits vaccinated with human chorionic gonadotropin ("HCG") revealed a prevalence of atherosclerotic lesions along the length of the aortas and particularly in the portion of the aortas from the thoracic region. In contrast, the aortas of rabbits vaccinated with a synthetic vaccine peptide having segments of tetanus toxoid and the C-terminal sequence of human CETP ("Peptide", see SEQ ID NO: 7), full-length recombinant human CETP ("rhuCETP"), and a CETP-tetanus toxoid conjugate composition ("Conjugate") had lower incidence of lesions, including the portion of the aorta from the thoracic region.

To quantitate the noticeable difference in the presence of atherosclerotic lesions in the aortas of rabbits or lack thereof, the total surface area of the pinned aortas and that of the aortic lesions was determined from photographs by planar morphometry (Daley et al., 1994) using a digitizing tablet with associated software (THE MORPHOMETER™, Woods Hole Educational Associates, Woods Hole, Massachusetts). The percentage of the surface area of the aortas covered by lesions was determined and the percentages are represented in Figure 11.

EXAMPLES 2 and 3

Plasmid-based Vaccines in Mice and Rabbits

Four groups of mice were vaccinated intramuscularly with one of the following:

1. pCMV: a plasmid vector having the cytomegalovirus immediate early promoter/enhancer but without any operably-linked structural gene
2. pCIII-huCETP: a plasmid vector having the full coding sequence for human CETP (SEQ ID NO: 1) under the transcriptional control of the human Apo CIII promoter
3. pSV40-huCETP: a plasmid vector having the full coding sequence for human CETP under the transcriptional control of the SV40 promoter

4. pCMV-TT-rabCETP: a plasmid vector having a tetanus toxoid peptide (amino acid : 2 to 15 of SEQ ID NO: 7) coding sequence and the full coding sequence for rabbit CETP (SEQ ID NO: 3) under the transcriptional control of the cytomegalovirus immediate early promoter/enhancer

5 The mice were injected once with 25 μ l of PBS containing 100 μ g of the plasmid and blood samples were periodically collected and were analyzed with an ELISA detecting anti-human CETP antibodies with the method described below:

Plastic 96-well microtiter plates were coated with Protein A/G, by incubating 100 μ l of a 5 μ g/ml PBS solution per well overnight at 4°C. The plates were emptied and the wells
10 were blocked with 200 μ l of blocking buffer (PBS with 4% BSA, 1% sucrose, 0.5% NP-40, 0.01% Gentamycin) for 2 to 8 hours at room temperature. Antibodies from the plasma samples were captured on the Protein A/G by incubating 100 μ l of various dilutions of the samples for 1 hour at room temperature. Following washing of the wells, biotinylated CETP was captured by the plate bound antibodies by incubating 100 μ l of a biotinylated CETP
15 solution at room temperature for 1 hour. The bound CETP was detected by incubating 100 μ l of a streptavidin-HRP solution for 30 minutes at room temperature, followed by adding 100 μ l of substrate, stopping the reaction with 0.18M sulfuric acid and reading the optical density at 450 nm. Figure 12 shows that plasmids delivered to Groups 2, 3, and 4 produce immunogenic xenogeneic protein.

20 Subsequently, rabbits were vaccinated with 300 μ g (equally split in six intramuscular sites in the quadriceps) of a vector carrying the human CETP coding sequence under the transcriptional control of the SV40 promoter enhancer (SV40-huCETP) or the same vector without the CETP coding sequence (SV40). The primary injection occurred on Day 1 with an identical boost on weeks 5, 8, 26, and 30. Blood samples were taken periodically throughout
25 the experiment and animals were terminated on week 34.

The plasma samples from the vaccinated rabbits were subjected to the ELISA for detection of antibodies to whole recombinant human CETP, as described above. Figure 13A-K summarizes the titration of the antibody measured in the rabbits vaccinated with the SV40 promoter enhancer (SV40-huCETP) between weeks 1-34. Significant
30 antibody production was detected on weeks 30 and 34, depicted by the open diamond and open square symbols of the graphs, in most animals. Two of the 11 rabbits (Figure 13A, and 13E) were non-responders.

The rabbits were switched from a diet of basic rabbit chow to diets supplemented with 0.25% cholesterol (w/w) known to produce atherosclerotic-like lesions in rabbits. The lesions in rabbits vaccinated with pSV40-huCETP and control rabbits vaccinated with pSV40 were visualized and quantitated as described in Example 1 above. Figure 14 shows the mean
5 percentage of aorta covered with lesions in both groups of rabbits. The results of this experiment are in line with the trend observed with directly vaccinated rabbits (Groups I-IV, see Figure 11), showing a decrease in aortic lesions due to vaccination.

Although a number of embodiments have been described above, it will be understood by those skilled in the art that modifications and variations of the described compositions and
10 methods may be made without departing from either the spirit of the invention or the scope of the appended claims. The articles and publications cited herein are incorporated by reference.

CLAIMS:

1. A vaccine composition effective to promote production of antibodies binding to endogenous cholesteryl ester transfer protein (CETP) in a mammal, consisting essentially of a non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.
2. A vaccine composition as defined in Claim 1, wherein said mammal is a human and said non-endogenous CETP is selected from the group consisting of rabbit CETP, mouse CETP, simian CETP, humanized rabbit, mouse or simian CETP, and allelic variants or polymorphs of said human's CETP.
3. A vaccine composition effective to promote production of antibodies binding to endogenous cholesteryl ester transfer protein (CETP) in a mammal, consisting essentially of a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.
4. A vaccine composition as defined in Claim 3, wherein said mammalianized CETP is a humanized rabbit CETP.
5. A vaccine composition as defined in Claim 4, wherein said humanized rabbit CETP has the amino acid sequence of SEQ ID NO. 5.
6. A vaccine composition as defined in Claim 1, wherein said mammal is a rabbit and said non-endogenous CETP is human CETP.
7. A vaccine composition as defined in Claim 1 or 3, which contains an adjuvant selected from the group consisting of alum, Freund's Complete Adjuvant, Freund's Incomplete Adjuvant, RIBI Adjuvant System.
8. A plasmid-based vaccine comprising a promoter sequence suitable for directing the transcription of a nucleotide sequence in a cell of a mammal operably linked to a nucleotide

sequence coding for a cholesteryl ester transfer protein (CETP) that is non-endogenous to said mammal.

9. A plasmid-based vaccine as defined in Claim 8, wherein said mammal is a human and said non-endogenous CETP is selected from the group consisting of rabbit CETP, mouse CETP, simian CETP, humanized rabbit, mouse or simian CETP, and allelic variants or polymorphs of said human's CETP.

10. A plasmid-based vaccine as defined in Claim 8, wherein said mammal is a human and said non-endogenous CETP is rabbit CETP.

11. A plasmid-based vaccine as defined in Claim 8, wherein said mammal is a human and said non-endogenous CETP is a humanized rabbit CETP.

12. A plasmid-based vaccine as defined in Claim 11, wherein said non-endogenous CETP has the amino acid sequence of SEQ ID NO: 5.

13. A plasmid-based vaccine as defined in Claim 8, wherein said mammal is a rabbit and said non-endogenous CETP is a human CETP.

14. A plasmid-based vaccine as defined in Claim 8, wherein the promoter is the cytomegalovirus immediate early promoter/enhancer.

15. A method for promoting production in a mammal of antibodies binding the mammal's endogenous CETP comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

16. A method for elevating the ratio of circulating high density lipoprotein-associated cholesterol to circulating low density lipoprotein-associated cholesterol or total cholesterol in a mammal comprising administering to the mammal a non-endogenous CETP or a

mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

17. A method for decreasing the level of endogenous CETP activity in a human or other animal comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

18. A method for altering the metabolism of low density lipoprotein-associated cholesterol to decrease the development of atherosclerotic lesions in a mammal comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

19. A method of lowering the level of circulating low density lipoprotein in a mammal comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

20. A method of lowering the level of total circulating cholesterol in a mammal comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

21. A method for therapeutically or prophylactically treating atherosclerosis in a mammal comprising administering to the mammal a non-endogenous CETP or a mammalianized non-endogenous CETP, optionally in combination with an adjuvant effective to non-specifically stimulate the immune response of said mammal.

22. A method according to any of Claims 15-21, wherein the mammal is a human and the non-endogenous CETP is a rabbit CETP.

23. A method according to any of Claims 15-21, wherein said mammalianized CETP is humanized rabbit CETP.
24. A method according to Claim 23, wherein said humanized rabbit CETP has the amino acid sequence of SEQ ID NO. 5.
25. A method according to any of Claims 15-21, wherein said CETP is administered in combination with an adjuvant selected from the group consisting of alum, Freund's Complete Adjuvant, Freund's Incomplete Adjuvant, RIBI Adjuvant System.
26. A method for promoting production in a mammal of antibodies reactive with the mammal's endogenous CETP comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
27. A method for elevating the ratio of circulating high density lipoprotein-associated cholesterol to circulating low density lipoprotein-associated cholesterol or total cholesterol in a mammal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
28. A method for decreasing the level of endogenous CETP activity in a human or other animal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
29. A method for altering the metabolism of low density lipoprotein-associated cholesterol to decrease the development of atherosclerotic lesions in a mammal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
30. A method of lowering the level of circulating low density lipoprotein in a mammal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.

31. A method of lowering the level of total circulating cholesterol in a mammal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
32. A method for therapeutically or prophylactically treating atherosclerosis in a mammal comprising transfecting at least one cell of the mammal with a plasmid-based vaccine according to Claim 8.
33. A method according to any of Claims 26-32, wherein the mammal is a human and the plasmid-based vaccine includes nucleic acid encoding a rabbit CETP.
34. A method according to any of Claims 26-32, wherein the mammal is a human and the plasmid-based vaccine includes nucleic acid encoding a humanized rabbit CETP.
35. A method according to Claims 34, wherein said humanized rabbit CETP has the amino acid sequence of SEQ ID NO. 5.
36. A method according to any of Claims 26-32, which comprises the additional step of administering to said mammal an adjuvant effective to non-specifically stimulate the immune response of said mammal.
37. A method according to Claim 36, wherein the adjuvant is selected from the group consisting of alum, Freund's Complete Adjuvant, Freund's Incomplete Adjuvant, RIBI Adjuvant System.

1/17

Cys	Pro	Lys	Gly	Ala	Ser	Tyr	Glu	Ala	Gly	Ile	Val	Cys	Arg	Ile	Thr	Lys	Pro	Ala	Leu	Leu	Val	Leu	Asn	Gln	25
Cys	Ser	Lys	Gly	Thr	Ser	His	Glu	Ala	Gly	Ile	Val	Cys	Arg	Ile	Thr	Lys	Pro	Ala	Leu	Leu	Val	Leu	Asn	His	
Glu	Thr	Ala	Lys	Val	Val	Gln	Thr	Ala	Phe	Gln	Arg	Ala	Gly	Tyr	Pro	Asp	Val	Ser	Gly	Glu	Arg	Ala	Val	Met	50
Glu	Thr	Ala	Lys	Val	Ile	Gln	Thr	Ala	Phe	Gln	Arg	Ala	Ser	Tyr	Pro	Asp	Ile	Thr	Gly	Glu	Lys	Ala	Met	Met	
Leu	Leu	Gly	Arg	Val	Lys	Tyr	Gly	Leu	His	Asn	Leu	Gln	Ile	Ser	His	Leu	Ser	Ile	Ala	Ser	Ser	Gln	Val	Glu	75
Leu	Leu	Gly	Gln	Val	Lys	Tyr	Gly	Leu	His	Asn	Ile	Gln	Ile	Ser	His	Leu	Ser	Ile	Ala	Ser	Ser	Gln	Val	Glu	
Leu	Val	Asp	Ala	Lys	Thr	Ile	Asp	Val	Ala	Ile	Gln	Asn	Val	Ser	Val	Val	Phe	Lys	Gly	Thr	Leu	Asn	Tyr	Ser	100
Leu	Val	Glu	Ala	Lys	Ser	Ile	Asp	Val	Ser	Ile	Gln	Asn	Val	Ser	Val	Val	Phe	Lys	Gly	Thr	Leu	Lys	Tyr	Gly	
Tyr	Thr	Ser	Ala	Trp	Gly	Leu	Gly	Ile	Asn	Gln	Ser	Val	Asp	Phe	Glu	Ile	Asp	Ser	Ala	Ile	Asp	Leu	Gln	Ile	125
Tyr	Thr	Thr	Ala	Trp	Trp	Leu	Gly	Ile	Asp	Gln	Ser	Ile	Asp	Phe	Glu	Ile	Asp	Ser	Ala	Ile	Asp	Leu	Gln	Ile	
Asn	Thr	Glu	Leu	Thr	Cys	Asp	Ala	Gly	Ser	Val	Arg	Thr	Asn	Ala	Pro	Asp	Cys	Tyr	Leu	Ala	Phe	His	Lys	Leu	150
Asn	Thr	Gln	Leu	Thr	Cys	Asp	Ser	Gly	Arg	Val	Arg	Thr	Asp	Ala	Pro	Asp	Cys	Tyr	Leu	Ser	Phe	His	Lys	Leu	
Leu	Leu	His	Leu	Gln	Gly	Glu	Arg	Glu	Pro	Gly	Trp	Leu	Lys	Gln	Leu	Phe	Thr	Asn	Phe	Ile	Ser	Phe	Thr	Leu	175
Leu	Leu	His	Leu	Gln	Gly	Glu	Arg	Glu	Pro	Gly	Trp	Ile	Lys	Gln	Leu	Phe	Thr	Asn	Phe	Ile	Ser	Phe	Thr	Leu	

FIG. 1A

2/17

Lys	Leu	Ile	Leu	Lys	Arg	Gln	Val	Cys	Asn	Glu	Ile	Asn	Thr	Ile	Ser	Asn	Ile	Met	Ala	Asp	Phe	Val	Gln	Thr	200
Lys	Leu	Val	Leu	Lys	Gly	Gln	Ile	Cys	Lys	Glu	Ile	Asn	Val	Ile	Ser	Asn	Ile	Met	Ala	Asp	Phe	Val	Gln	Thr	
Arg	Ala	Ala	Ser	Ile	Leu	Ser	Asp	Gly	Asp	Ile	Gly	Val	Asp	Ile	Ser	Val	Thr	Gly	Ala	Pro	Val	Ile	Thr	Ala	225
Arg	Ala	Ala	Ser	Ile	Leu	Ser	Asp	Gly	Asp	Ile	Gly	Val	Asp	Ile	Ser	Leu	Thr	Gly	Asp	Pro	Val	Ile	Thr	Ala	
Thr	Tyr	Leu	Glu	Ser	His	His	Lys	Gly	His	Phe	Thr	His	Lys	Asn	Val	Ser	Glu	Ala	Phe	Pro	Leu	Arg	Ala	Phe	250
Ser	Tyr	Leu	Glu	Ser	His	His	Lys	Gly	His	Phe	Ile	Tyr	Lys	Asn	Val	Ser	Glu	Asp	Leu	Pro	Leu	Pro	Thr	Phe	
Pro	Pro	Gly	Leu	Leu	Gly	Asp	Ser	Arg	Met	Leu	Tyr	Phe	Thr	Phe	Ser	Asp	Gln	Val	Leu	Asn	Ser	Leu	Ala	Arg	275
Ser	Pro	Thr	Leu	Leu	Gly	Asp	Ser	Arg	Met	Leu	Tyr	Phe	Thr	Phe	Ser	Glu	Arg	Val	Phe	His	Ser	Leu	Ala	Lys	
Ala	Ala	Phe	Gln	Glu	Gly	Arg	Leu	Val	Leu	Ser	Leu	Thr	Gly	Asp	Glu	Phe	Lys	Lys	Val	Leu	Glu	Thr	Gln	Gly	300
Val	Ala	Phe	Gln	Asp	Gly	Arg	Leu	Met	Leu	Ser	Leu	Met	Gly	Asp	Glu	Phe	Lys	Ala	Val	Leu	Glu	Thr	Trp	Gly	
Phe	Asp	Thr	Asn	Gln	Glu	Ile	Phe	Gln	Glu	Leu	Ser	Arg	Gly	Leu	Pro	Thr	Gly	Gln	Ala	Gln	Val	Ala	Val	His	325
Phe	Asn	Thr	Asn	Gln	Glu	Ile	Phe	Gln	Glu	val	val	Gly	Gly	Phe	Pro	Ser	-----Gln	Ala	Gln	val	Thr	Val	His		
Cys	Leu	Lys	Val	Pro	Lys	Ile	Ser	Cys	Gln	Asn	Arg	Gly	val	val	Val	Ser	Ser	Ser	Val	Ala	val	Thr	Phe	Arg	350
Cys	Leu	Lys	Met	Pro	Lys	Ile	Ser	Cys	Gln	Asn	Lys	Gly	val	val	Val	Asn	Ser	Ser	Val	Met	val	Lys	Phe	Leu	

FIG. 1B

3/17

```

Phe Pro Arg Pro Asp Gly Arg Glu Ala Val Ala Tyr Arg Phe Glu Glu Asp Ile Ile Thr Thr Val Gln Ala Ser 375
| | | | | | | | | | | | | | | | | | | | | |
Phe Pro Arg Pro Asp Gln Gln His Ser Val Ala Tyr Thr Phe Glu Glu Asp Ile Val Thr Thr Val Gln Val Ser |
| | | | | | | | | | | | | | | | | | | | | |

Tyr Ser Gln Lys Lys Leu Phe Leu His Leu Leu Asp Phe Gln Cys Val Pro Ala Ser Gly Arg Ala Gly Ser Ser 400
| | | | | | | | | | | | | | | | | | | | | |
Tyr Ser Lys Lys Lys Leu Phe Leu Ser Leu Leu Asp Phe Gln Ile Thr Pro-----
| | | | | | | | | | | | | | | | | | | | | |

Ala Asn Leu Ser Val Ala Leu Arg Thr Glu Ala Lys Ala Val Ser Asn Leu Thr Glu Ser Arg Ser Glu Ser Leu 425
| | | | | | | | | | | | | | | | | | | | | |
-----Lys Thr Val Ser Asn Leu Thr Glu Ser Ser Ser Glu Ser Ile
| | | | | | | | | | | | | | | | | | | | | |

Gln Ser Ser Leu Arg Ser Leu Ile Ala Thr Val Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Ala Phe Thr 450
| | | | | | | | | | | | | | | | | | | | | |
Gln Ser Phe Leu Gln Ser Met Ile Thr Ala Val Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Val Phe Thr
| | | | | | | | | | | | | | | | | | | | | |

Ala Leu Met Asn Ser Lys Gly Leu Asp Leu Phe Glu Ile Ile Asn Pro Glu Ile Ile Thr Leu Asp Gly Cys Leu 475
| | | | | | | | | | | | | | | | | | | | | |
Ala Leu Met Asn Ser Lys Gly Val Ser Leu Phe Asp Ile Ile Asn Pro Glu Ile Ile Thr Arg Asp Gly Phe Leu
| | | | | | | | | | | | | | | | | | | | | |

Leu Leu Gln Met Asp Phe Gly Phe Pro Lys His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser (SEQ ID NO: 3) 496
| | | | | | | | | | | | | | | | | | | | | |
Leu Leu Gln Met Asp Phe Gly Phe Pro Glu His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser (SEQ ID NO: 1)
| | | | | | | | | | | | | | | | | | | | | |

```

FIG. 1C

4/17




FIG. 2

5/17




FIG. 3

6/17




FIG. 4

7/17




FIG. 5

8/17




FIG. 6

9/17




FIG. 7

9/17




FIG. 7

10/17




FIG. 8

11/17




FIG. 9

12/17




FIG. 10

13/17




FIG. 11

14/17




FIG. 12

15/17




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 13D




FIG. 13E





FIG. 13F



16/17




FIG. 13G




FIG. 13H




FIG. 13I




FIG. 13J





FIG. 13K



17/17




FIG. 14

SEQUENCE LISTING

<110> Rittershaus, Charles W.
 Thomas, Lawrence J.
 Avant Immunotherapeutics, Inc.

<120> Xenogeneic Cholesteryl Ester Transfer Protein (CETP)
 for Modulation of CETP Activity

<130> TCS-420.1 PCT seqlist

<140> PCT/US98/22145

<141> 1998-10-20

<150> 08/954,643

<151> 1997-10-20

<160> 7

<170> PatentIn Ver. 2.0

<210> 1

<211> 476

<212> PRT

<213> Homo sapiens

<300>

<301> Drayna, Dennis

<302> Cloning and Sequencing of Human Cholesteryl Ester
 Transfer cDNA

<303> Nature

<304> 327

<306> 632-634

<307> 1987-06-18

<313> 1 TO 476

<400> 1

Cys Ser Lys Gly Thr Ser His Glu Ala Gly Ile Val Cys Arg Ile Thr
 1 5 10 15

Lys Pro Ala Leu Leu Val Leu Asn His Glu Thr Ala Lys Val Ile Gln
 20 25 30

Thr Ala Phe Gln Arg Ala Ser Tyr Pro Asp Ile Thr Gly Glu Lys Ala
 35 40 45

Met Met Leu Leu Gly Gln Val Lys Tyr Gly Leu His Asn Ile Gln Ile
 50 55 60

Ser His Leu Ser Ile Ala Ser Ser Gln Val Glu Leu Val Glu Ala Lys
 65 70 75 80

Ser Ile Asp Val Ser Ile Gln Asn Val Ser Val Val Phe Lys Gly Thr
 85 90 95

Leu Lys Tyr Gly Tyr Thr Thr Ala Trp Trp Leu Gly Ile Asp Gln Ser
 100 105 110

Ile Asp Phe Glu Ile Asp Ser Ala Ile Asp Leu Gln Ile Asn Thr Gln
 115 120 125

Leu Thr Cys Asp Ser Gly Arg Val Arg Thr Asp Ala Pro Asp Cys Tyr
 130 135 140

Leu Ser Phe His Lys Leu Leu Leu His Leu Gln Gly Glu Arg Glu Pro
 145 150 155 160

Gly Trp Ile Lys Gln Leu Phe Thr Asn Phe Ile Ser Phe Thr Leu Lys
 165 170 175

Leu Val Leu Lys Gly Gln Ile Cys Lys Glu Ile Asn Val Ile Ser Asn
 180 185 190

Ile Met Ala Asp Phe Val Gln Thr Arg Ala Ala Ser Ile Leu Ser Asp
 195 200 205

Gly Asp Ile Gly Val Asp Ile Ser Leu Thr Gly Asp Pro Val Ile Thr
 210 215 220

Ala Ser Tyr Leu Glu Ser His His Lys Gly His Phe Ile Tyr Lys Asn
 225 230 235 240

Val Ser Glu Asp Leu Pro Leu Pro Thr Phe Ser Pro Thr Leu Leu Gly
 245 250 255

Asp Ser Arg Met Leu Tyr Phe Trp Phe Ser Glu Arg Val Phe His Ser
 260 265 270

Leu Ala Lys Val Ala Phe Gln Asp Gly Arg Leu Met Leu Ser Leu Met
 275 280 285

Gly Asp Glu Phe Lys Ala Val Leu Glu Thr Trp Gly Phe Asn Thr Asn
 290 295 300

Gln Glu Ile Phe Gln Glu Val Val Gly Gly Phe Pro Ser Gln Ala Gln
 305 310 315 320

Val Thr Val His Cys Leu Lys Met Pro Lys Ile Ser Cys Gln Asn Lys
 325 330 335

Gly Val Val Val Asn Ser Ser Val Met Val Lys Phe Leu Phe Pro Arg
 340 345 350

Pro Asp Gln Gln His Ser Val Ala Tyr Thr Phe Glu Glu Asp Ile Val
 355 360 365

Thr Thr Val Gln Ala Ser Tyr Ser Lys Lys Lys Leu Phe Leu Ser Leu
 370 375 380

Leu Asp Phe Gln Ile Thr Pro Lys Thr Val Ser Asn Leu Thr Glu Ser
 385 390 395 400

Ser Ser Glu Ser Ile Gln Ser Phe Leu Gln Ser Met Ile Thr Ala Val
 405 410 415

Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Val Phe Thr Ala Leu
 420 425 430

Met Asn Ser Lys Gly Val Ser Leu Phe Asp Ile Ile Asn Pro Glu Ile
 435 440 445

Ile Thr Arg Asp Gly Phe Leu Leu Leu Gln Met Asp Phe Gly Phe Pro
 450 455 460

Glu His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser
 465 470 475

<210> 2

<211> 1428

<212> DNA

<213> Homo sapiens

<300>

<301> Drayna, Dennis

<302> Cloning and Sequencing of Human Cholesteryl Ester
 Transfer cDNA

<303> Nature

<304> 327

<306> 632-634

<307> 1987-06-18

<313> 1 TO 476

<400> 2

```

tgctccaaag gcacctcgca cgaggcagge atcgtgtgcc gcatcaccaa gcctgccctc 60
ctgggtgttg accacgagac tgccaagggtg atccagaccg ccttcagcg agccagctac 120
ccagatatca cgggcgagaa ggccatgatg ctccctggcc aagtcaagta tgggttgac 180
aacatccaga tcagccaatt gtccatcgcc agcagccagg tggagctggt ggaagccaag 240
tccattgatg tctccattca gaacgtgtct gtgggtcttca aggggaccct gaagtatggc 300
tacaccactg cctgggtggct ggggtattgat cagtccattg acttcgagat cgactctgcc 360
attgacctcc agatcaaacac acagctgacc tgtgactctg gtagagtgcg gaccgatgcc 420
cctgactgct acctgtcttt ccataagctg ctccctgcac tccaagggga gcgagagcct 480
gggtggatca agcagctggt cacaaatttc atctccttca ccctgaagct ggtcctgaag 540
ggacagatct gcaaagagat caacgtcatc tctaacatca tggccgattt tgtccagaca 600
agggtgacca gcatcctttc agatggagac attgggggtg acatttccct gacaggtgat 660
cccgtcatca cagcctccta cctggagtcc catcacaagg gtcatttcat ctacaagaat 720
gtctcagagg acctccccct cccacacctc tcgcccacac tgctggggga ctcccgcacg 780
ctgtacttct ggttctctga gcgagtcttc cactcgctgg ccaaggtagc tttccaggat 840
ggccgcctca tgctcagcct gatgggagac gagttcaagg cagtgtctga gacctggggc 900
ttcaaacacca accaggaat cttccaagag gttgtcggcg gcttccccag ccaggcccaa 960
gtcaccgtec actgcctcaa gatgcccaag atctcctgcc aaaacaaggg agtcgtggtc 1020
aattcttcag tgatgggtgaa attcctcttt ccacgcccag accagcaaca tctgttagct 1080
tacacatttg aagaggatat cgtgactacc gtccaggcct cctattctaa gaaaaagctc 1140
ttcttaagcc tcttgattt ccagattaca ccaaagactg tttccaactt gactgagagc 1200
agctccgagt ccatccagag ctccctgcag tcaatgatca ccgtgtggg catccctgag 1260
gtcatgtctc ggctcgaggt agtgtttaca gccctcatga acagcaaagg cgtgagcctc 1320
ttcgacatca tcaaccctga gattatcact cgagatggct tcctgctgct gcagatggac 1380
tttggtctcc ctgagcacct gctggtggat ttcctccaga gcttgagc 1428

```

<210> 3

<211> 496

<212> PRT

<213> *Oryctolagus cuniculus*

<300>

<301> Nagashima, Mariko

<302> Cloning and mRNA tissue distribution of rabbit
cholesteryl ester transfer protein

<303> J. Lipid Res.

<304> 29

<306> 1643-1649

<307> 1988

<313> 1 TO 496

<400> 3

Cys Pro Lys Gly Ala Ser Tyr Glu Ala Gly Ile Val Cys Arg Ile Thr

1

5

10

15

Lys Pro Ala Leu Leu Val Leu Asn Gln Glu Thr Ala Lys Val Val Gln

20

25

30

Thr Ala Phe Gln Arg Ala Gly Tyr Pro Asp Val Ser Gly Glu Arg Ala

35	40	45
Val Met Leu Leu Gly Arg Val Lys Tyr Gly Leu His Asn Leu Gln Ile		
50	55	60
Ser His Leu Ser Ile Ala Ser Ser Gln Val Glu Leu Val Asp Ala Lys		
65	70	75 80
Thr Ile Asp Val Ala Ile Gln Asn Val Ser Val Val Phe Lys Gly Thr		
85	90	95
Leu Asn Tyr Ser Tyr Thr Ser Ala Trp Gly Leu Gly Ile Asn Gln Ser		
100	105	110
Val Asp Phe Glu Ile Asp Ser Ala Ile Asp Leu Gln Ile Asn Thr Glu		
115	120	125
Leu Thr Cys Asp Ala Gly Ser Val Arg Thr Asn Ala Pro Asp Cys Tyr		
130	135	140
Leu Ala Phe His Lys Leu Leu Leu His Leu Gln Gly Glu Arg Glu Pro		
145	150	155 160
Gly Trp Leu Lys Gln Leu Phe Thr Asn Phe Ile Ser Phe Thr Leu Lys		
165	170	175
Leu Ile Leu Lys Arg Gln Val Cys Asn Glu Ile Asn Thr Ile Ser Asn		
180	185	190
Ile Met Ala Asp Phe Val Gln Thr Arg Ala Ala Ser Ile Leu Ser Asp		
195	200	205
Gly Asp Ile Gly Val Asp Ile Ser Val Thr Gly Ala Pro Val Ile Thr		
210	215	220
Ala Thr Tyr Leu Glu Ser His His Lys Gly His Phe Thr His Lys Asn		
225	230	235 240
Val Ser Glu Ala Phe Pro Leu Arg Ala Phe Pro Pro Gly Leu Leu Gly		
245	250	255
Asp Ser Arg Met Leu Tyr Phe Trp Phe Ser Asp Gln Val Leu Asn Ser		
260	265	270
Leu Ala Arg Ala Ala Phe Gln Glu Gly Arg Leu Val Leu Ser Leu Thr		
275	280	285
Gly Asp Glu Phe Lys Lys Val Leu Glu Thr Gln Gly Phe Asp Thr Asn		

290	295	300
Gln Glu Ile Phe Gln Glu Leu Ser Arg Gly Leu Pro Thr Gly Gln Ala		
305	310	315 320
Gln Val Ala Val His Cys Leu Lys Val Pro Lys Ile Ser Cys Gln Asn		
	325	330 335
Arg Gly Val Val Val Ser Ser Ser Val Ala Val Thr Phe Arg Phe Pro		
	340	345 350
Arg Pro Asp Gly Arg Glu Ala Val Ala Tyr Arg Phe Glu Glu Asp Ile		
	355	360 365
Ile Thr Thr Val Gln Ala Ser Tyr Ser Gln Lys Lys Leu Phe Leu His		
	370	375 380
Leu Leu Asp Phe Gln Cys Val Pro Ala Ser Gly Arg Ala Gly Ser Ser		
385	390	395 400
Ala Asn Leu Ser Val Ala Leu Arg Thr Glu Ala Lys Ala Val Ser Asn		
	405	410 415
Leu Thr Glu Ser Arg Ser Glu Ser Leu Gln Ser Ser Leu Arg Ser Leu		
	420	425 430
Ile Ala Thr Val Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Ala		
	435	440 445
Phe Thr Ala Leu Met Asn Ser Lys Gly Leu Asp Leu Phe Glu Ile Ile		
	450	455 460
Asn Pro Glu Ile Ile Thr Leu Asp Gly Cys Leu Leu Leu Gln Met Asp		
465	470	475 480
Phe Gly Phe Pro Lys His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser		
	485	490 495

<210> 4

<211> 1488

<212> DNA

<213> Oryctolagus cuniculus

<300>

<301> Nagashima, Mariko
 <302> Cloning and mRNA tissue distribution of rabbit
 cholesteryl ester transfer protein
 <303> J. Lipid Res.
 <304> 29
 <306> 1643-1649
 <307> 1988
 <313> 1 TO 496

<400> 4
 tgtcccaag ggcctccta cgaggctggc atcgtgtgtc gcatcaccaa gcccgccctc 60
 ttggtgttga accaagagac ggccaagggtg gtccagacgg ccttccagcg cgccggctat 120
 ccggacgtca ggcgcgagag ggccgtgatg ctccctcgcc gggtaagta cgggctgcac 180
 aacctccaga tcagccacct gtccatcgcc agcagccagg tggagctggg ggacgccaaag 240
 accatcgacg tcgccatcca gaacgtgtcc gtggtcttca aggggaccct gaactacagc 300
 tacacgagtg cctgggggtt gggcatcaat cagtctgtcg acttcgagat cgactctgcc 360
 attgacctcc agatcaaacac agagctgacc tgcgacgtg gcagtgtgcg caccaatgcc 420
 cccgactgct acctggcttt ccataaactg ctccctgcacc tccaggggga ggcgcgagccg 480
 ggggtggctca agcagctctt cacaaacttc atctccttca ccctgaagct gattctgaag 540
 cgacaggtct gcaatgagat caacaccatc tccaacatca tggctgactt tgtccagacg 600
 agggccgcca gcatcctctc agatggagac atcgggggtg acatttccgt gacggggggc 660
 cctgtcatca cagccaccta cctggagtcc catcacaagg gtcacttcac gcacaagaac 720
 gtctccgagg ccttccccct ccgcgccttc ccgcccgtc ttctggggga ctcccgcatg 780
 ctctacttct ggttctccga tcaagtgtc aactccctgg ccagggccgc cttccaggag 840
 ggccgtctcg tgtcagcct gacaggggat gagttcaaga aagtgtgga gaccaggggt 900
 ttcgacacca accaggaaat cttccaggag ctttccagag gccttcccac cggccaggcc 960
 caggtagccg tccactgcct taagggtgcc aagatctcct gccagaaccg ggggtgtcgtg 1020
 gtgtcttctt ccgtcgccgt gacgttccgc tccccccg cagatggccg agaagctgtg 1080
 gcctacaggt ttgaggagga tatcatcacc accgtccagg cctcctactc ccagaaaaag 1140
 ctcttctctac acctcttga tttccagtgc gtgccggcca gcggaagggc aggcagctca 1200
 gcaaatctct ccgtggccct caggactgag gctaaggctg tttccaacct gactgagagc 1260
 cgctccgagt ccttcgagag ctctctccgc tccctgatcg ccacggtggg catcccggag 1320
 gtcattgtct ggctcgaggt ggcgttcaca gccctcatga acagcaaagg cctggacctc 1380
 ttcgaaatca tcaaccccg gattatcact ctgatggct gcctgctgct gcagatggac 1440
 ttcgggtttt ccaagcacct gctgggtggat ttctgcaga gcctgagc 1488

<210> 5
 <211> 477
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: humanized
 rabbit CETP protein

<400> 5
 Cys Pro Lys Gly Ala Ser Tyr Glu Ala Gly Ile Val Cys Arg Ile Thr
 1 5 10 15

Lys Pro Ala Leu Leu Val Leu Asn Gln Glu Thr Ala Lys Val Val Gln
 20 25 30

Thr Ala Phe Gln Arg Ala Gly Tyr Pro Asp Val Ser Gly Glu Arg Ala
 35 40 45

Val Met Leu Leu Gly Arg Val Lys Tyr Gly Leu His Asn Leu Gln Ile
 50 55 60

Ser His Leu Ser Ile Ala Ser Ser Gln Val Glu Leu Val Asp Ala Lys
 65 70 75 80

Thr Ile Asp Val Ala Ile Gln Asn Val Ser Val Val Phe Lys Gly Thr
 85 90 95

Leu Asn Tyr Ser Tyr Thr Ser Ala Trp Gly Leu Gly Ile Asn Gln Ser
 100 105 110

Val Asp Phe Glu Ile Asp Ser Ala Ile Asp Leu Gln Ile Asn Thr Glu
 115 120 125

Leu Thr Cys Asp Ala Gly Ser Val Arg Thr Asn Ala Pro Asp Cys Tyr
 130 135 140

Leu Ala Phe His Lys Leu Leu Leu His Leu Gln Gly Glu Arg Glu Pro
 145 150 155 160

Gly Trp Leu Lys Gln Leu Phe Thr Asn Phe Ile Ser Phe Thr Leu Lys
 165 170 175

Leu Ile Leu Lys Arg Gln Val Cys Asn Glu Ile Asn Thr Ile Ser Asn
 180 185 190

Ile Met Ala Asp Phe Val Gln Thr Arg Ala Ala Ser Ile Leu Ser Asp
 195 200 205

Gly Asp Ile Gly Val Asp Ile Ser Val Thr Gly Ala Pro Val Ile Thr
 210 215 220

Ala Thr Tyr Leu Glu Ser His His Lys Gly His Phe Thr His Lys Asn
 225 230 235 240

Val Ser Glu Ala Phe Pro Leu Arg Ala Phe Pro Pro Gly Leu Leu Gly
 245 250 255

Asp Ser Arg Met Leu Tyr Phe Trp Phe Ser Asp Gln Val Leu Asn Ser
 260 265 270

Leu Ala Arg Ala Ala Phe Gln Glu Gly Arg Leu Val Leu Ser Leu Thr
 275 280 285

Gly Asp Glu Phe Lys Lys Val Leu Glu Thr Gln Gly Phe Asp Thr Asn
 290 295 300

Gln Glu Ile Phe Gln Glu Leu Ser Arg Gly Leu Pro Thr Gly Gln Ala
 305 310 315 320

Gln Val Ala Val His Cys Leu Lys Val Pro Lys Ile Ser Cys Gln Asn
 325 330 335

Arg Gly Val Val Val Ser Ser Ser Val Ala Val Thr Phe Arg Phe Pro
 340 345 350

Arg Pro Asp Gly Arg Glu Ala Val Ala Tyr Arg Phe Glu Glu Asp Ile
 355 360 365

Ile Thr Thr Val Gln Ala Ser Tyr Ser Gln Lys Lys Leu Phe Leu His
 370 375 380

Leu Leu Asp Phe Gln Cys Val Pro Lys Ala Val Ser Asn Leu Thr Glu
 385 390 395 400

Ser Arg Ser Glu Ser Leu Gln Ser Ser Leu Arg Ser Leu Ile Ala Thr
 405 410 415

Val Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Ala Phe Thr Ala
 420 425 430

Leu Met Asn Ser Lys Gly Leu Asp Leu Phe Glu Ile Ile Asn Pro Glu
 435 440 445

Ile Ile Thr Leu Asp Gly Cys Leu Leu Leu Gln Met Asp Phe Gly Phe
 450 455 460

Pro Lys His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser
 465 470 475

<210> 6

<211> 496

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: humanized

rabbit CESTP protein

<400> 6

Cys Pro Lys Gly Ala Ser Tyr Glu Ala Gly Ile Val Cys Arg Ile Thr
 1 5 10 15
 Lys Pro Ala Leu Leu Val Leu Asn Gln Glu Thr Ala Lys Val Val Gln
 20 25 30
 Thr Ala Phe Gln Arg Ala Gly Tyr Pro Asp Val Ser Gly Glu Arg Ala
 35 40 45
 Val Met Leu Leu Gly Arg Val Lys Tyr Gly Leu His Asn Leu Gln Ile
 50 55 60
 Ser His Leu Ser Ile Ala Ser Ser Gln Val Glu Leu Val Asp Ala Lys
 65 70 75 80
 Thr Ile Asp Val Ala Ile Gln Asn Val Ser Val Val Phe Lys Gly Thr
 85 90 95
 Leu Asn Tyr Ser Tyr Thr Ser Ala Trp Gly Leu Gly Ile Asn Gln Ser
 100 105 110
 Val Asp Phe Glu Ile Asp Ser Ala Ile Asp Leu Gln Ile Asn Thr Glu
 115 120 125
 Leu Thr Cys Asp Ala Gly Ser Val Arg Thr Asn Ala Pro Asp Cys Tyr
 130 135 140
 Leu Ala Phe His Lys Leu Leu Leu His Leu Gln Gly Glu Arg Glu Pro
 145 150 155 160
 Gly Trp Leu Lys Gln Leu Phe Thr Asn Phe Ile Ser Phe Thr Leu Lys
 165 170 175
 Leu Ile Leu Lys Arg Gln Val Cys Asn Glu Ile Asn Thr Ile Ser Asn
 180 185 190
 Ile Met Ala Asp Phe Val Gln Thr Arg Ala Ala Ser Ile Leu Ser Asp
 195 200 205
 Gly Asp Ile Gly Val Asp Ile Ser Val Thr Gly Ala Pro Val Ile Thr
 210 215 220
 Ala Thr Tyr Leu Glu Ser His His Lys Gly His Phe Thr His Lys Asn
 225 230 235 240

Val Ser Glu Ala Phe Pro Leu Arg Ala Phe Pro Pro Gly Leu Leu Gly
 245 250 255
 Asp Ser Arg Met Leu Tyr Phe Trp Phe Ser Asp Gln Val Leu Asn Ser
 260 265 270
 Leu Ala Arg Ala Ala Phe Gln Glu Gly Arg Leu Val Leu Ser Leu Thr
 275 280 285
 Gly Asp Glu Phe Lys Lys Val Leu Glu Thr Gln Gly Phe Asp Thr Asn
 290 295 300
 Gln Glu Ile Phe Gln Glu Leu Ser Arg Gly Leu Pro Thr Gly Gln Ala
 305 310 315 320
 Gln Val Ala Val His Cys Leu Lys Val Pro Lys Ile Ser Cys Gln Asn
 325 330 335
 Arg Gly Val Val Val Ser Ser Ser Val Ala Val Thr Phe Arg Phe Pro
 340 345 350
 Arg Pro Asp Gly Arg Glu Ala Val Ala Tyr Arg Phe Glu Glu Asp Ile
 355 360 365
 Ile Thr Thr Val Gln Ala Ser Tyr Ser Gln Lys Lys Leu Phe Leu His
 370 375 380
 Leu Leu Asp Phe Gln Cys Val Pro Ala Ser Gly Arg Ala Gly Ser Ser
 385 390 395 400
 Ala Asn Leu Ser Val Ala Leu Arg Thr Glu Ala Lys Ala Val Ser Asn
 405 410 415
 Leu Thr Glu Ser Arg Ser Glu Ser Leu Gln Ser Ser Leu Arg Ser Leu
 420 425 430
 Ile Ala Thr Val Gly Ile Pro Glu Val Met Ser Arg Leu Glu Val Ala
 435 440 445
 Phe Thr Ala Leu Met Asn Ser Lys Gly Leu Asp Leu Phe Glu Ile Ile
 450 455 460
 Asn Pro Glu Ile Ile Thr Leu Asp Gly Cys Leu Leu Leu Gln Met Asp
 465 470 475 480
 Phe Gly Phe Pro Glu His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser
 485 490 495

<210> 7

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: fusion protein
of a tetanus toxoid segment and human C-terminus
C-terminus

<400> 7

Cys Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu Phe
1 5 10 15

Gly Phe Pro Glu His Leu Leu Val Asp Phe Leu Gln Ser Leu Ser
20 25 30

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/22145

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K39/00 A61K39/39 A61K48/00 C07K14/47

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HESLER C. ET AL.: "Purification and characterization of a human plasma cholesteryl ester transfer protein" J. BIOL. CHEM, vol. 262, no. 5, - 15 February 1987 pages 2275-2282, XP002096997 see whole doc., esp. P.2282 2.col.	1,3,6,15
X	SMITH A M ET AL: "PREPARATION OF AN ANTI-PEPTIDE ANTISERUM SPECIFIC FOR CHOLESTERYL ESTER TRANSFER PROTEIN (CEPT)" MEDICAL SCIENCE RESEARCH, vol. 21, no. 24, 16 December 1993, page 911/912 XP002041707 see the whole document -/-	1,3,7,15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

18 March 1999

Date of mailing of the international search report

01/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Müller, F

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/22145

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 39168 A (IMMUNE RESPONSE CORP INC) 12 December 1996 SEE WHOLE DOC., esp. page 1,2 and claims ---	1-3,7, 15-21
X	WO 96 34888 A (T CELL SCIENCES INC ;RITTERSHAUS CHARLES W (US); THOMAS LAWRENCE J) 7 November 1996 cited in the application see whole doc. esp. claims and examples ---	1-3,6,7, 15-17, 20,21
A	HESLER C. ET AL.,: "Monoclonal antibodies to the Mr 74000 cholesteryl ester transfer protein neutralize all of the cholesteryl ester and triglyceride transfer activities in human plasma" J. BIOL. CHEM., vol. 263, no. 11, - 15 April 1988 pages 5020-5023, XP002096998 ---	1-37
P,X	WO 97 41227 A (T CELL SCIENCES INC ;THOMAS LAWRENCE J (US)) 6 November 1997 cited in the application see the whole document ---	8-14, 26-37
P,X	THOMAS L.J. ET AL.,: "Use of xenogeneic cholesteryl ester transfer protein (CETP) in a plasmid-based vaccine to produce anti-cetp autoantibodies for the prevention/treatment of atherosclerosis" FASEB JOURNAL, vol. 12, no. 4, - 17 March 1998 page a310,1805 XP002096999 see the whole document -----	8-14, 26-37

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/22145

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 15-37 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internati Application No

PCT/US 98/22145

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9639168 A	12-12-1996	AU 6091296 A CA 2223177 A EP 0831881 A	24-12-1996 12-12-1996 01-04-1998
WO 9634888 A	07-11-1996	AU 5636096 A CA 2219795 A EP 0827509 A	21-11-1996 07-11-1996 11-03-1998
WO 9741227 A	06-11-1997	AU 2994697 A	19-11-1997